



DUX4 gene

double homeobox 4

Normal Function

The *DUX4* gene is located near the end of chromosome 4 in a region known as D4Z4. This region consists of 11 to more than 100 repeated segments, each of which is about 3,300 DNA base pairs (3.3 kb) long. The entire D4Z4 region is normally hypermethylated, which means that it has a large number of methyl groups (consisting of one carbon atom and three hydrogen atoms) attached to the DNA. The addition of methyl groups turns off (silences) genes, so hypermethylated regions of DNA tend to have fewer genes that are turned on (active).

Each of the repeated segments in the D4Z4 region contains a copy of the *DUX4* gene; the copy closest to the end of chromosome 4 is called *DUX4*, while the other copies are described as "*DUX4*-like" or *DUX4L*. Hypermethylation of the D4Z4 region keeps the *DUX4*-like genes silenced all the time. No protein is produced from these genes. The *DUX4* gene is also silenced in most adult cells and tissues, although it is active during early development and in the testes of adult males. Little is known about the function of the protein produced from the active *DUX4* gene; it appears to help control the activity of other genes.

The *DUX4* gene (the copy closest to the end of chromosome 4) is located next to a regulatory region of DNA known as a pLAM sequence, which is necessary for the production of the DUX4 protein. Some copies of chromosome 4 have a functional pLAM sequence, while others do not. Copies of chromosome 4 with a functional pLAM sequence are described as 4qA or "permissive." Those without a functional pLAM sequence are described as 4qB or "non-permissive." Without a functional pLAM sequence, no DUX4 protein is made. Because there are two copies of chromosome 4 in each cell, individuals may have two "permissive" copies of chromosome 4, two "non-permissive" copies, or one of each.

Health Conditions Related to Genetic Changes

facioscapulohumeral muscular dystrophy

Changes in the D4Z4 region of chromosome 4, which contains the *DUX4* gene, cause facioscapulohumeral muscular dystrophy. This disorder is characterized by muscle weakness and wasting (atrophy) that worsens slowly over time. Two types of the disorder have been described: type 1 (FSHD1) and type 2 (FSHD2). Both types result from hypomethylation of the D4Z4 region, in which the DNA has fewer methyl groups attached than normal. In FSHD1, hypomethylation occurs because the D4Z4


region is abnormally shortened (contracted), containing between 1 and 10 repeats instead of the usual 11 to 100 repeats. In FSHD2, hypomethylation most often results from mutations in a gene called *SMCHD1*, which normally hypermethylates the D4Z4 region.

Hypomethylation of the D4Z4 region prevents the *DUX4* gene from being silenced in cells and tissues where it is usually turned off, such as adult muscle cells. However, hypomethylation of the D4Z4 region results in facioscapulohumeral muscular dystrophy only when it occurs with a "permissive" chromosome 4. The "permissive" chromosome contains a working pLAM sequence, which allows protein to be produced from the abnormally active *DUX4* gene. Researchers believe that the protein influences the activity of other genes, particularly in muscle cells. However, it is unknown how presence of the DUX4 protein damages or destroys these cells, leading to progressive muscle weakness and atrophy.

Chromosomal Location

Cytogenetic Location: 4q35.2, which is the long (q) arm of chromosome 4 at position 35.2

Molecular Location: base pairs 190,173,701 to 190,175,845 on chromosome 4 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- double homeobox 4-like
- double homeobox protein 4
- double homeobox protein 4/10
- double homeobox protein 10

Additional Information & Resources

Educational Resources

- Developmental Biology (sixth edition, 2000): Methylation Pattern and the Control of Transcription
<https://www.ncbi.nlm.nih.gov/books/NBK10038/>

GeneReviews

- Facioscapulohumeral Muscular Dystrophy
<https://www.ncbi.nlm.nih.gov/books/NBK1443>

Scientific Articles on PubMed

- PubMed
https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28DUX4%5BTIAB%5D%29+OR+%28double+homeobox+4%5BTIAB%5D%29+OR+%28DUX4L*%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D

OMIM

- DOUBLE HOMEBOX PROTEIN 4
<http://omim.org/entry/606009>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_DUX4.html
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=50800
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/100288687>
- UniProt
<http://www.uniprot.org/uniprot/Q9UBX2>

Sources for This Summary

- OMIM: DOUBLE HOMEBOX PROTEIN 4
<http://omim.org/entry/606009>
- Geng LN, Yao Z, Snider L, Fong AP, Cech JN, Young JM, van der Maarel SM, Ruzzo WL, Gentleman RC, Tawil R, Tapscott SJ. DUX4 activates germline genes, retroelements, and immune mediators: implications for facioscapulohumeral dystrophy. *Dev Cell*. 2012 Jan 17;22(1):38-51. doi: 10.1016/j.devcel.2011.11.013. Epub 2011 Dec 29.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22209328>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3264808/>

- Lemmers RJ, Tawil R, Petek LM, Balog J, Block GJ, Santen GW, Amell AM, van der Vliet PJ, Almomani R, Straasheijm KR, Krom YD, Klooster R, Sun Y, den Dunnen JT, Helmer Q, Donlin-Smith CM, Padberg GW, van Engelen BG, de Greef JC, Aartsma-Rus AM, Frants RR, de Visser M, Desnuelle C, Sacconi S, Filippova GN, Bakker B, Bamshad MJ, Tapscott SJ, Miller DG, van der Maarel SM. Digenic inheritance of an SMCHD1 mutation and an FSHD-permissive D4Z4 allele causes facioscapulohumeral muscular dystrophy type 2. *Nat Genet.* 2012 Dec;44(12):1370-4. doi: 10.1038/ng.2454. Epub 2012 Nov 11.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/23143600>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3671095/>
- Lemmers RJ, van der Vliet PJ, Klooster R, Sacconi S, Camaño P, Dauwerse JG, Snider L, Straasheijm KR, van Ommen GJ, Padberg GW, Miller DG, Tapscott SJ, Tawil R, Frants RR, van der Maarel SM. A unifying genetic model for facioscapulohumeral muscular dystrophy. *Science.* 2010 Sep 24;329(5999):1650-3. doi: 10.1126/science.1189044. Epub 2010 Aug 19.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/20724583>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4677822/>
- Sacconi S, Salviati L, Desnuelle C. Facioscapulohumeral muscular dystrophy. *Biochim Biophys Acta.* 2015 Apr;1852(4):607-14. doi: 10.1016/j.bbadis.2014.05.021. Epub 2014 May 29. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24882751>
- Tawil R, van der Maarel SM, Tapscott SJ. Facioscapulohumeral dystrophy: the path to consensus on pathophysiology. *Skelet Muscle.* 2014 Jun 10;4:12. doi: 10.1186/2044-5040-4-12. eCollection 2014. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24940479>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4060068/>
- Yao Z, Snider L, Balog J, Lemmers RJ, Van Der Maarel SM, Tawil R, Tapscott SJ. DUX4-induced gene expression is the major molecular signature in FSHD skeletal muscle. *Hum Mol Genet.* 2014 Oct 15;23(20):5342-52. doi: 10.1093/hmg/ddu251. Epub 2014 May 26.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24861551>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4168822/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/DUX4>

Reviewed: August 2014
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services